

Results and Discussion

Only alcohol non-drinkers (27.1% v 19.4%; $p=0.01$) and current smoking (44.6% v 52.9%; $p=0.02$) differed significantly between NSWPHS and NSW-NATSIHS (Table 1).

The demographic characteristics of the NSWPHS and NSW-NATSIHS samples were also compared to the 2001 Aboriginal and Torres Strait Islander population of NSW.⁸ The NSWPHS sample contained a higher proportion of older adults, females, and regional and remote adults, and the NSW-NATSIHS sample contained a higher proportion of females, and regional and remote adults.

Finding that the majority of the risk behaviours and all the health status prevalence estimates were not significantly different was encouraging. The higher rate of alcohol non-drinking and lower rate of current smoking in the NSWPHS was likely due to the under-representation of young Aboriginal and Torres Strait Islander men in the NSWPHS telephone survey. Although 98% of Aboriginal and Torres Strait Islander persons had used telephones in the month prior to the National Aboriginal and Torres Strait Islander Social Survey (NATSISS) interview, landline phone ownership still remains low (71% in non-remote and 40% remote) compared to the general public; however, mobile phone ownership is becoming more common (more than 90% in non-remote and more than 80% remote).⁹ Expanding the NSWPHS sampling frame, as planned in 2012, to include mobile phones could potentially increase access to young Aboriginal and Torres Strait Islander men and improve the accuracy of the estimates for alcohol and smoking.

References

1. Australian Bureau of Statistics. 3302.0.55.003 – *Experimental Life Tables for Aboriginal and Torres Strait Islander Australians, 2005–2007*. Canberra (AUST): ABS; 2009 May.
2. Australian Institute of Health and Welfare. *Aboriginal and Torres Strait Islander Health Performance Framework 2010: Detailed Analyses*. Catalogue No.: IHW 53. Canberra (AUST): AIHW; 2011 Sept.
3. Australian Bureau of Statistics. 4715.0. – *National Aboriginal and Torres Strait Islander Health Survey, 2004-05*. Canberra (AUST): ABS; 2006 April.
4. NSW Ministry of Health. *Adult Population Health Survey* [Internet]. Sydney (AUST): State Government of New South Wales; 2010 [cited 2012 Jul 30]. Available from: <http://www.health.nsw.gov.au/publichealth/surveys/index.asp>
5. Centre for Epidemiology and Research. *Report on Adult Aboriginal Health from the New South Wales Population Health Survey 2002-2005*. Sydney (AUST): NSW Department of Health; 2006 Jan.
6. Centre for Epidemiology and Research. *Report on Adult Aboriginal Health from the New South Wales Population Health Survey 2006-2009*. Sydney (AUST): NSW Department of Health; 2010 Nov.
7. Australian Bureau of Statistics. 4713.0. – *Population Characteristics, Aboriginal and Torres Strait Islander Australians, 2001*. Canberra (AUST): ABS; 2003 Oct.
8. Altman DG, Bland JM. Interaction revisited: the difference between two estimates. *BMJ*. 2003;326:219.
9. Australian Bureau of Statistics. 4714.0. – *National Aboriginal and Torres Strait Islander Social Survey, 2008*. Canberra (AUST): ABS; 2009 Oct.

Correspondence to: Ms Margo Barr, Centre for Epidemiology and Evidence, NSW Ministry of Health, 73 Miller St, North Sydney, NSW 2055; e-mail: margo.barr@doh.health.nsw.gov.au

doi: 10.1111/1753-6405.12018

Associations between discrimination and dental visiting behaviours in an Aboriginal Australian birth cohort

Lisa M. Jamieson, Margie Steffens

Australian Research Centre for Population Oral Health, The University of Adelaide, South Australia

Yin C. Paradies

McCaughy Centre, University of Melbourne, Victoria

Race-based discrimination is associated with a range of adverse health conditions among Indigenous Australians. Studies show links between race-based discrimination, and depression and anxiety as well as smoking, substance use, psychological distress and poor self-assessed health status.¹ The Australian Medical Association reported evidence of ‘inherent discrimination’ in the Australian health system, finding the medical attention received by Indigenous Australians is frequently ‘culturally intolerant’ and unwelcoming.²

There is limited information regarding the role of discrimination in access to dental services among Indigenous Australians. We explored whether (i) there were associations between dental service access and self-reported discrimination and (ii) if yes, whether associations between discrimination and access to dental services persisted after adjustment for other risk factors. Participants were Wave-3 members of the Aboriginal Birth Cohort study, a prospective longitudinal investigation of Indigenous Australians living in the Northern Territory’s Top End. Participants completed a self-report dental questionnaire. The item pertaining to use of dental services was ‘Have you seen a dentist before?’, with response options of ‘yes’ and ‘no’. Participants additionally completed a social and emotional well-being instrument. Based on items used in the National Aboriginal and Torres Strait Islander Health Survey, respondents were asked if they had ever been treated unfairly or discriminated against because they were Aboriginal. Response options were dichotomised into ‘not really or little bit’ and ‘fair bit and lots of times’. Other risk factors included sex, age, household size, oral health impairment and dental fear. Explanatory variables, which were all significant at the $P<0.1$ level in bivariate analysis, were entered into a logistical regression model. Adjusted odds ratios were considered statistically significant when p -values derived from the Wald statistic were ≤ 0.05 . Data were analysed using PASW version 18.

Of the 468 participants for whom vital status was obtained, 442 provided complete information in the self-report dental questionnaire, which was 95% of the total number of participants examined in Wave-3 and 69% of those recruited at birth (i.e. Wave-1) who were still alive. Data on self-reported discrimination were available for 336 (76%) of those individuals and all subsequent analyses were limited to this sample. There were no significant dental utilisation differences between those included and not included in the analysis. Eighteen participants (5.4%) reported having never visited a dentist before (Table 1) while 31 (9.2%) reported race-based discrimination. In bivariate analyses, the only significant association with having never visited a dentist before was self-reported discrimination. This

finding persisted after adjusting for other risk factors in multivariate modeling, with those reporting discrimination being 3.8 times more likely to have never visited a dentist before.

The results of this exploratory study highlight the importance of examining discrimination-related factors when determining the broad range of barriers associated with poor Indigenous dental service utilisation rates. The findings additionally support inclusion of discrimination awareness and anti-discrimination initiatives when developing interventions aimed at fostering improved uptake in dental service utilisation among Indigenous Australian groups.

The study has a number of limitations. The outcome of 'not visited' was quite uncommon at 5.4%, as was the discrimination variable at 9.2% of respondents. While this reduces the external validity of the study (i.e. its population representativeness), it does not limit the internal validity; small sample sizes are a problem because they make statistically significant findings much more unlikely. When significant differences do occur, as with our study, the findings are arguably more convincing as such an association must be strong to be significant in such a small sample. It is also important to note that a single item does not capture the full extent of discrimination or its impact on health and health service utilisation.

Although more sophisticated designs among larger cohorts are necessary to explore these findings further, it is notable that discrimination was the only significant risk factor and that this persisted after adjustment in multivariate modelling. International

research suggests that experiencing race-based discrimination in healthcare settings leads to disengagement with, and avoidance of, healthcare in the future.³ Participants' views in a qualitative study suggest that formation of a strong relationship between oral health service providers and Indigenous clients is paramount to the utilisation of, and satisfaction with, these services for Indigenous clients.⁴

In summary, the findings from this exploratory research suggest that race-based discrimination may play a role in the under-utilisation of dental services among Australian Aboriginal young adults. Further research with larger samples is required to more clearly elucidate the causal pathways between these outcomes.

References

1. Paradies Y, Harris R, Anderson I. *The Impact of Racism on Indigenous Health in Australia and Aotearoa: Towards a Research Agenda, Discussion Paper No. 4*. Darwin (AUST): Cooperative Research Centre for Aboriginal Health; 2008.
2. Australian Medical Association. *Institutionalised Inequity Not Just a Matter of Money*. Canberra (AUST): AMA; 2007.
3. Lee C, Ayers SL, Kronenfeld JJ. The association between perceived provider discrimination, healthcare utilization and health status in racial and ethnic minorities. *Ethn Dis*. 2009;193:330-7.
4. Jamieson LM, Parker EJ, Richards L. Using qualitative methodology to inform an Indigenous-owned oral health promotion initiative in Australia. *Health Promot Int*. 2008;23:52-9.

Correspondence to: Lisa M. Jamieson, Australian Research Centre for Population Oral Health, The University of Adelaide, South Australia 5005, e-mail: lisa.jamieson@adelaide.edu.au

Table 1: Dental visiting behaviours of ABC study participants by dental-related factors.

	Number of participants (%)	Number not visited dentist before (%)	Unadjusted OR (95% CI)	Adjusted OR ^c (95% CI)
Total	336 (100.0)	18 (5.4)		
Socio-demographic				
Sex				
Male	156 (46.4)	7 (4.5)	1	1
Female	180 (53.6)	11 (6.1)	1.39 (0.52–3.67)	1.42 (0.50–4.05)
Age				
16-18 years	234 (69.6)	15 (6.4)	1	1
19-20 years	102 (30.4)	3 (2.9)	0.44 (0.13–1.56)	0.44 (0.12–1.60)
Household size				
Four or less people	70 (20.8)	1 (1.4)	1	1
Five or more people	247 (73.5)	14 (4.8)	4.78 (0.62–36.69)	4.39 (0.56–34.56)
Oral health impairment^b				
No	226 (67.3)	15 (6.6)	1	1
Yes	110 (32.7)	3 (2.7)	2.54 (0.72–8.95)	2.49 (0.66–9.34)
Would you feel scared about going to the dentist?				
No	138 (41.1)	4 (2.9)	1	1
Little bit, fair bit, heaps	198 (58.9)	14 (7.1)	2.55 (0.82–7.92)	1.70 (0.52–5.61)
Have you been treated unfairly or discriminated against because you are Aboriginal?				
Not really or little bit	305 (90.8)	13 (4.3)	1	1
Fair bit or lots	31 (9.2)	5 (16.1) ^a	4.32 (1.43–13.06) ^a	3.79 (1.18–12.06) ^a

Notes:

^a $p < 0.05$

^b 'Do you have any trouble with your teeth, gum or jaw right now?' OR 'Since the last Wet, have you stopped eating some foods because they hurt your teeth?' OR a response of 'Some or none good' to the item 'Do you think your teeth are looking ok?'

^c Adjusting for all other covariates in the model