

Birth weight and cognitive function in early adulthood: the Australian aboriginal birth cohort study

M. S. Pearce^{1*}, K. D. Mann¹, G. Singh² and S. M. Sayers²

¹*Institute of Health & Society, Newcastle University, UK*

²*Menzies School of Health Research, Charles Darwin University, Darwin, Australia*

It has been suggested that in addition to genetic factors, fetal and post-natal growth influence cognition in early adulthood. However, most studies have been in developed populations, so it is unclear if the same findings would be seen in other, less developed, settings, and have used testing tools not applicable to an Australia Aboriginal population. This study investigated the relationships between cognitive function in early adulthood and birth weight and contemporary height. Simple reaction time (SRT), choice reaction time (CRT) and working memory (WM) were assessed using the CogState battery. A significant association was seen between birth weight and SRT in early adulthood, but not with the other two cognitive measures. Urban dwellers had significantly shorter SRT and CRT than their remote counterparts. Contemporary body mass index and maternal age were associated with CRT. Only fetal growth restriction was associated with WM, with greater WM in those with restricted growth. No associations were seen with contemporary height. These results suggest that fetal growth may be more important than the factors influencing post-natal growth in terms of cognition in early adulthood in this population, but that the associations may be inconsistent between cognitive outcomes. Further research is required to identify whether similar associations are seen in other, similar, populations and to assess why differences in cognitive outcome measures are seen.

Received 22 November 2013; Revised 7 January 2014; Accepted 9 January 2014; First published online 14 February 2014

Key words: birth cohort, birth weight, cognition, Indigenous populations

Introduction

Although cognitive ability has a heritable component, environmental influences are also likely to play a substantial role. Nutritional status *in utero*, for which birth weight is often used as an indicator, is one such environmental influence that may affect brain development.¹ However, as birth weight also has a considerable heritable component, any association with birth weight may be equally mediated by genetic factors.²

Very low birth weight babies are known to be less well developed in terms of cognitive function in childhood,^{3–6} and early adulthood.^{5,7–10} An association between birth weight and later cognition, even within the normal birth weight range in term births, has been seen in some studies,^{11–15} but not in others.^{16–18}

Height in adulthood has been consistently related to cognition in adulthood.^{7,19–21} Less is known regarding the relationship between cognition and height in early adulthood, although height in childhood has been shown to be positively associated with cognitive ability.^{7,17,22} However, it is unclear whether such a relationship persists, or whether it diminishes with age. Obesity has been associated with cognitive function at a range of ages,²³ although the association may have a bi-directional component, particularly in later life. Childhood malnutrition has also been linked to poorer cognition.²⁴

The vast majority of the studies linking either birth weight or height with cognition have been in developed countries, so it is unclear if the same findings would be seen in other, less developed, settings. Although Australia is a developed country, it includes a disadvantaged Aboriginal Australian population. Australian Aborigines have low birth weight rates more than double those of the non-Aboriginal population (13% compared with 6%).²⁵ With such high rates of adverse fetal growth, it is important to study whether the links between birth weight and later cognition are seen in this population. Moreover, traditional cognitive assessment tools are mostly inappropriate for populations such as Indigenous Australians, as they tend to rely heavily on the use of both spoken and written English language and unfamiliar concepts.

This study, using prospectively recorded data from the Australian Aboriginal birth cohort study,²⁶ investigated the relationships between cognitive function in early adulthood and birth weight and contemporary height.

Materials and methods

Participants in this study were members of the Australian Aboriginal birth cohort. To be eligible for this cohort, a baby had to be a singleton born at Royal Darwin Hospital between January 1987 and March 1990 to a mother self-identified and recorded as Aboriginal in the Delivery Suite Register. This resulted in 686 Aboriginal babies (of the 1238 eligible babies) being recruited.²⁶ There were no significant differences in the mean birth weight, low birth weight rates or sex ratio between

*Address for correspondence: M. S. Pearce, Institute of Health & Society, Newcastle University, Newcastle upon Tyne, NE1 4LP, UK.
(Email mark.pearce@ncl.ac.uk)

those recruited and those not.²⁶ At the time of recruitment, the Royal Darwin Hospital was the routine place of delivery for 98% of Aboriginal mothers within the local health region of 120,000 km², and the tertiary referral hospital for high risk deliveries transferred *in utero* from a sparsely populated vast area covering 2 million km² of northern Australia. Hence the cohort consists of both routine deliveries and also those deliveries resulting from the high risk *in uteri* referrals.

Information on a number of factors, including birth weight, gestational age, maternal age, growth, residential status (as a surrogate for socio-economic status), and cognition was recorded prospectively for study members. Measures of weight, length and head circumference were taken at birth. Gestational age was estimated within 4 days of birth by the same neonatal paediatrician using the Dubowitz scoring system,²⁷ previously evaluated for Aboriginal babies.²⁸

Along with the continuous measure, birth weight was categorized into two groups for size; <2.5 and >2.5 kg and a further two groups for fetal growth restriction; <10th percentile or not, using post-natal gestational age estimations and an Australian-based reference standard contemporary with the time of recruitment.²⁹

Residence, at the time of follow up, was defined as remote (residence in defined remote Aboriginal communities) or non-remote (residence within the twin cities of Darwin and Palmerston, the greater Darwin area and smaller rural towns).

Measures of adult height, weight, age and cognition were taken between December 2005 and January 2008. Participants were measured while wearing light clothing and no shoes. Weight was measured, once, to the nearest 0.1 kg using a digital electronic scale (model TBF-521; Tanita Corp, Arlington Heights, IL), and height to the nearest millimetre with a portable wall-mounted stadiometer. Body mass index (BMI) was calculated from height and weight. Height from this period was standardized using the Centers for Disease Control and Prevention (CDC) 2000 growth reference.³⁰ Standardised training was given to all assessors of height and weight. Maternal age was collected from the hospital birth records at the time of recruitment of the baby into the study.

Cognitive assessment

Cognitive function was assessed using the CogState computerized cognitive test battery (CogState Ltd, Melbourne, Australia, www.cogstate.com). CogState is a non-verbal, computerized, cognitive assessment developed for the assessment of diverse groups,³¹ and previously shown to be valid in terms of reliability and minimal practice effects in a sample of Indigenous Australian adolescents.³² The battery in this study consisted of three tasks that are based on playing cards displayed on a computer screen. By using playing cards, it is independent of language, culture and socio-economic background. The tasks were as follows:

(1) Simple reaction time (SRT): This is a measure of psychomotor speed in which one playing card is shown

face down on the computer screen, with an instruction to press the 'yes' key as quickly as possible whenever the card is turned face up. This is repeated 35 times, randomly. A lower score reflects better performance.

(2) Choice reaction time (CRT): This measures CRT and decision making attention. A face-down playing card is displayed on the screen. When the card is flipped over, if it is red, the participant should press the 'yes' key, and if it is not red, then the participant should press the 'no' key. This is repeated 30 times, again randomly. A lower score reflects better performance.

(3) WM: This measures primarily WM, but also psychomotor speed and visual attention. Each time a card is revealed, the participant must decide whether he/she has been shown that card before in this task and respond by pressing the 'yes' or 'no' key. The participant should, therefore, try to remember all the cards that are presented in this task. The procedure is repeated 30 times, again randomly. A higher score reflects better performance.

The tasks within the CogState battery were always tested in the order in which they are listed above. The overall duration for the battery to be completed was between 7 and 10 min.

Statistical analysis

Sex differences in cognition (SRT, CRT, WM) were tested using the Wilcoxon rank sum test. Associations between cognition and explanatory variables were estimated using linear regression. Regression coefficients are presented with accompanying 95% confidence intervals. Gestational age, birth weight, age, heights, BMI and maternal age were treated as continuous variables. Residential status sex, low birth weight and fetal growth restriction were treated as categorical variables. Cohort members with missing data were included in all analyses for which they contributed complete data (i.e. complete data for all variables included in a particular model). Multiple adjusted models were formed to ensure that the various measures of birth size and fetal growth were not included in the same models. All statistical analyses were done using the Stata statistical software package, version 12.0 (StataCorp, College Station, TX).

Results

During data collection between 2005 and 2008, 263, 255 and 259 study members took part in the SRT, CRT and WM tasks, respectively (Table 1). Of those that took part in the cognition testing, 46% were male (54% female) and 74% were remote residents. Despite small differences in the participation numbers between the different cognitive tests, means and standard deviations/medians and inter-quartile ranges were the same across variables investigated, therefore descriptive statistics are shown for the maximum sample available in Table 1.

No difference in any of the cognitive outcome variables (SRT, CRT and WM) was seen between males and females

Table 1. Descriptive statistics, continuous variables

Variable	<i>n</i>	Mean (SD)/median (IQR)
Birth weight (kg)	263	3.00 (0.61)
Gestational age (weeks)	237	38 (1.84)
Age (years)	262	18 (1.09)
Height at current age (cm)	262	167.69 (8.91)
Height at current age (<i>Z</i> -score)	262	-0.20 (0.91)
BMI at current age (<i>Z</i> -score)	262	-0.46 (1.63)
Maternal age (years)	262	22 (5.78)
SRT	263	345.74 (283.35, 468.52)
CRT	255	613.90 (517.23, 722.78)
WM	259	1089.53 (853.29, 1386.00)

IQR, inter-quartile range; BMI, body mass index; SRT, simple reaction time; CRT, choice reaction time; WM, working memory.

($P = 0.424, 0.826, 0.552$, respectively). A significant negative association between SRT and birth weight was seen (Table 2) which remained, with greater magnitude, after adjustment for gestational age, maternal age and residential status (Table 3). This significant negative association also remained when birth weight was categorized, above and below 2.5 kg ($P = 0.046$, adjusted $P = 0.023$) with low birth weight associated with longer reaction times. There were neither significant associations between gestational age and cognition, nor any association with birth weight and the remaining tests. Having had fetal growth restriction was significantly associated with longer SRT ($P = 0.002$, adjusted $P = 0.008$). However, having had fetal growth restriction was also associated with a greater proportion of correct responses to the WM task (i.e. better WM) ($P = 0.014$, adjusted $P = 0.023$). There were 28 individuals born pre-term (i.e. <37 weeks gestation) that completed the CogState battery. Being born premature was not significantly associated with any of the cognitive tests.

Age of the participants undertaking cognitive assessments ranged from 17 to 19 years (Table 1) and no significant associations were seen between age and any of the outcomes. Maternal age, at the date of birth of the participants, ranged from 18 to 26 years and also showed no significant associations with any of the outcomes.

Both contemporary height measures (raw and *Z*-score) showed significant, negative, associations with both SRT and CRT (Table 2). However, neither remained significant after adjustment for gestational age, maternal age and residential status (Table 3). BMI (*Z*-score) was significantly negatively associated with SRT and CRT, but not with WM. After adjustment for gestational age, maternal age and residential status, the remaining association with CRT showed a decrease in reaction time with increasing BMI *Z*-score.

Those with non-remote residential status performed significantly better in SRT and CRT than those with remote residence. While a borderline significant result with WM (with a reduced proportion of correct responses in the WM task

among urban participants), this was not significant in the adjusted model.

Models were valid in terms of adhering to the assumptions underlying linear regression and no outliers were detected.

Discussion

In this cohort of Indigenous Australians, a significant association was seen between birth weight and SRT in early adulthood, but not with the other two cognitive measures. Urban residents had significantly lower SRT and CRT than their remote counterparts. Contemporary BMI and maternal age were associated with CRT, while only fetal growth restriction was associated with WM. No associations were seen with contemporary height.

Our finding of a negative association between birth weight and SRT is consistent with previous studies that have shown associations between low birth weight and cognition in early adulthood.^{5,7-15} Specifically to reaction times, Strang-Karlsson *et al.*'s study of the Helsinki birth cohort showed that adults who were born at very low birth weights had slower reaction times than control adults.⁹ However, while our findings are consistent for SRT this appears to be the first time this has been shown in the whole birth weight range of a population, Strang-Karlsson *et al.* also showed associations with other cognitive outcomes from the CogState battery, including CRT and WM, with worse performance in those born at very low birth weights.⁹ A study of neuromotor function also found lower SRT in children (5-7 years old) who had been very low birth weight babies, although this was a cycling-based test, rather than the card-based test used in this study.³³ The finding of greater WM in those with fetal growth restriction was in the opposite direction to that expected, and to that reported by Strang-Karlsson *et al.*⁹ While this may reflect a difference in the study populations included, or relate to an unmeasured exposure or compensatory mechanism during the prenatal or post-natal period, it may also be due to residual confounding or chance.

While previous studies have shown associations between height in adulthood and cognition in adulthood^{7,19-21} and between height in childhood and cognitive ability,^{7,17,22} no such associations were seen in this study. No previous associations appear to have been reported between adult height and any of the three specific cognitive outcomes in this study.

A number of previous studies have shown a stronger effect of socio-economic status on childhood cognition than that seen for birth weight.^{16,34-35} There is no measure of socio-economic status available for the Australian Aboriginal population. However, the residential status of remote and non-remote recorded at the time of follow-up, can be used as proxy for socio-economic status with the presumption of being more disadvantaged with a residential status of remote. Those with a non-remote residence had lower simple and CRT when compared with those in a remote residence. This may reflect such factors as schooling and exposure to computers which are

Table 2. Univariate linear regression results

Variable		Co-efficient	95% CI	P	n
Gestational age (weeks)					
SRT		3.22	-10.55, 16.99	0.645	237
CRT		2.73	-9.19, 14.65	0.652	229
WM		29.48	-32.67, 91.63	0.351	233
Birth weight (kg)					
SRT		-61.01	-100.04, -21.97	0.002	263
CRT		-24.03	-58.86, 10.80	0.125	255
WM		-56.81	-244.26, 130.65	0.551	259
Birth weight (categorized)					
SRT	>2.5 kg	Reference			263
	<2.5 kg	68.51	1.29, 135.73	0.046	
CRT	>2.5 kg	Reference			255
	<2.5 kg	46.71	-12.48, 105.91	0.121	
WM	>2.5 kg	Reference			259
	<2.5 kg	155.45	-163.30, 474.20	0.338	
Fetal growth restriction, birth weight <10th percentile					
SRT	No	Reference			237
	Yes	87.44	32.03, 142.85	0.002	
CRT	No	Reference			229
	Yes	28.47	-20.79, 77.72	0.256	
WM	No	Reference			233
	Yes	317.85	63.79, 571.91	0.014	
Age (years)					
SRT		4.52	-17.57, 26.61	0.687	262
CRT		3.77	-15.73, 23.26	0.704	254
WM		14.7	-89.90, 119.30	0.782	258
Height at current age (cm)					
SRT		-2.94	-5.63, -0.25	0.032	262
CRT		-2.41	-4.78, -0.03	0.047	254
WM		8.84	-3.94, 21.6	0.174	258
Height at current age (Z-score)					
SRT		-39.06	-65.23, -12.87	0.004	262
CRT		-35.12	-58.26, -11.98	0.003	254
WM		29.15	-96.70, 154.99	0.649	258
BMI at current age (Z-score)					
SRT		-28.68	-43.09, -14.28	<0.001	262
CRT		-18.79	-31.77, -5.80	0.005	254
WM		-48.67	-119.07, 21.73	0.175	258
Region					
SRT	Remote	Reference			262
	Urban	-144.32	-197.07, -91.58	<0.001	
CRT	Remote	Reference			254
	Non-remote	-123.26	-169.72, -76.81	<0.001	
WM	Remote	Reference			258
	Non-remote	-237.49	-498.52, 23.54	0.074	
Maternal age (years)					
SRT		-1.21	-5.40, 2.98	0.570	262
CRT		2.1	-1.57, 5.78	0.261	254
WM		-2.32	-22.17, 17.52	0.818	258

SRT, simple reaction time; CRT, choice reaction time; WM, working memory.

greater with a non-remote residence and less likely to reflect nutritional status which would be expected to operate in the opposite direction.

Obesity has been associated with cognitive function at a range of ages,²³ although the association may have a bi-directional component. In contrast, we found a negative

Table 3. Results of the adjusted linear regression models for the three cognitive outcomes

Variable	Co-efficient	95% CI	P
SRT (n = 235)			
Gestational age (weeks)	18.15	1.41, 34.89	0.034
Birth weight (kg) ^a	-76.39	-128.16, -24.63	0.004
Birth weight (categorized) ^a			
> 2.5 kg	Reference		
< 2.5 kg	103.02	14.15, 191.87	0.023
Fetal growth restriction, birth weight <10th percentile ^a			
No	Reference		
Yes	72.97	19.09, 126.86	0.008
Age (years)	0.91	-20.86, 22.67	0.934
Height at current age (cm)	-0.58	-3.65, 2.50	0.712
Height at current age (Z-score)	-14.68	-44.39, 15.02	0.331
BMI at current age (Z-score)	-12.45	-28.71, 3.82	0.133
Residential status			
Remote	Reference		
Non-remote	-130.96	-189.17, -72.74	<0.001
Maternal age (years)	1.92	-2.40, 6.24	0.382
CRT (n = 227)			
Gestational age (weeks)	5.07	-9.62, 19.76	0.497
Birth weight (kg) ^a	-16.31	-61.63, 29.01	0.479
Birth weight (categorized) ^a			
> 2.5 kg	Reference		
< 2.5 kg	53.52	-23.39, 130.43	0.172
Fetal growth restriction, birth weight <10th percentile ^a			
No	Reference		
Yes	6.97	-55.85, 69.79	0.827
Age (years)	3.89	-15.13, 22.90	0.687
Height at current age (cm)	-1.64	-4.34, 1.05	0.232
Height at current age (Z-score)	-22.29	-48.50, 3.92	0.095
BMI at current age (Z-score)	-14.38	-28.67, -0.11	0.048
Residential status			
Remote	Reference		
Non-remote	-122.06	-172.96, -71.16	<0.001
Maternal age (years)	4.19	0.40, 7.97	0.030
WM (n = 231)			
Gestational age (weeks)	59.13	-21.23, 139.49	0.148
Birth weight (kg) ^a	-142.31	-390.01, 105.39	0.259
Birth weight (categorized) ^a			
> 2.5 kg	Reference		
< 2.5 kg	405.18	-14.07, 824.44	0.058
Fetal growth restriction, birth weight <10th percentile ^a			
No	Reference		
Yes	297.56	40.48, 554.64	0.023
Age (years)	28.57	-75.73, 132.87	0.590
Height at current age (cm)	7.12	-7.64, 21.89	0.343
Height at current age (Z-score)	2.06	-141.39, 145.51	0.977
BMI at current age (Z-score)	20.5	-58.26, 99.27	0.608
Residential status			
Remote	Reference		
Non-remote	-199.18	-477.73, 79.38	0.160
Maternal age (years)	6.78	-13.97, 27.53	0.521

SRT, simple reaction time; BMI, body mass index; CRT, choice reaction time; WM, working memory.

^aBirth weight was not included as an adjustment for birth size or fetal growth-related variables. These early growth data were also not included within the same models, with only birth weight (kg) used as an adjustment factor for the non-fetal growth related variables.

association between contemporary BMI and CRT. Given that the mean BMI *Z*-score was nearly half a standard deviation below zero, this does not reflect a likely influence of obesity or overweight, of which there were few in this population. Instead, better cognitive performance in terms of CRT in those with higher BMI is likely to reflect nutritional status being better in those in the normal BMI range, and that underweight can be associated with cognition as well as overweight.²⁴

Strengths and weaknesses

The main strength of this study is that prospectively collected data on early life experience, including fetal growth, gestational age and maternal age, could be analysed, in relation to three cognitive outcomes variables, alongside later measures of adult height and residential status in an Australian Aboriginal population. Rather than using the cognitive assessment methods often used in populations in developed countries, we used the CogState battery, which is much more appropriate and less prone to bias in this population. To our knowledge, no other study has used the CogState battery to assess associations between early growth and later cognition over the entire birth weight range.

It has also been proposed that genetic factors and parental education levels may mediate the relationship between birth weight and cognitive ability.^{2,36} Maternal education and duration of breastfeeding have been linked with cognition.³⁷ We were unable to take account of these possible confounding factors in this investigation. However, for breastfeeding, as the vast majority of Aboriginal infants were breastfed,³⁸ this is unlikely to have introduced much in the way of confounding. Similarly data were not available for maternal height or BMI.

Although a postnatal classification of fetal growth restriction may not be ideal, it was necessary in this study as 70% of mothers did not know their last menstrual period and only 7% of mothers had a dating ultrasound before 14 weeks.²⁸ The Dubowitz scoring system used has been evaluated previously and, was found to provide valid estimates of gestational age in Aboriginal neonates.²⁸ The use of the CDC growth reference fits the CDC view that only set of growth charts is needed to cover all racial and ethnic groups and is supported by research evidence from a number of studies, including one in the Canadian Aboriginal population.³⁹ While the high breastfeeding rates in the Aboriginal population may make using the CDC references difficult in early life, this is less of an issue for the heights measured at the same as the cognitive assessments.

We were not able to control for the presence of colour-blindness or other problems with vision or with fine motor skills. However, there are likely to be few participants with conditions severe enough to impact on performance in the CogState battery, and such individuals would be unlikely to take part in the battery (or give a very low score, which we did not see). We were also unable to assess changes in nutritional status in childhood and how that would relate to changes in neurodevelopmental outcomes.

Different findings in terms of associations with the three cognitive outcome variables may reflect an issue with the sample size in this population, in that additional significant findings may have been seen with a larger sample. However, it may also reflect true differences in which factors influence different aspects of cognition, or differences between the way factors influence cognition in this population compared with other populations. In particular, the inconsistent findings for early growth are mirrored by the findings for residential status, with both fetal growth restriction and remote residence associated with greater WM.

Conclusion

In this cohort of Indigenous Australians, significant associations were seen between a range of factors and cognition in early adulthood. However, the associations differed depending on the outcome measure, but included birth weight inversely associated with SRT and fetal growth restriction associated with greater WM. Reaction times differed between non-remote and remote-based participants, suggesting that aspects of life, possibly related to socio-economic status play a role in influencing this aspect of cognitive function. No associations were seen with contemporary height. These results suggest that fetal growth may be more important than the factors influencing post-natal growth in terms of cognition in early adulthood in this population, but may be inconsistent between different cognitive outcomes. Further research is required to identify whether similar associations are seen in other, similar, populations, and to assess why differences in cognitive outcome measures are seen.

Acknowledgements

The authors thank the dedicated ABC research team who traced participants and collected the data. They thank Dr Sheree Cairney for helping the study to access and use the CogState battery. They also thank the young adults belonging to the cohort for their co-operation and all the individuals who helped in the urban and rural locations.

Financial Support

This work was supported by the National Health and Medical Research Council of Australia, the Channel 7 Children's Research Foundation of South Australia, the National Heart Foundation and a Northern Territory Government Research and Innovation Grant. MSP and KDM's collaboration with this study was funded by a Royal Society Travel Exchange Award.

Conflicts of Interest

None.

Ethical Standards

The Human Research Ethics Committee of the NT Department of Health & Families and Menzies School of Health Research, including the Aboriginal Ethical Sub-committee which has the power of veto, approved the study. Written consent was obtained in the form of an itemized consent with participant allowed to refuse individual procedures.

References

- Morgane PJ, Austin-La France R, Bronzino J, et al. Prenatal malnutrition and development of the brain. *Neurosci Biobehav Rev.* 1993; 17, 91–128.
- Boomsma DI, van Beijsterveldt CEM, Rietveld MJH, Bartels M, van Baal GC. Genetics mediate relation of birth weight to childhood IQ. *Br Med J.* 2001; 323, 1426–1427.
- Drillien CM. The incidence of mental and physical handicaps in school age children of very low birth weight. II. *Pediatrics.* 1967; 39, 238–247.
- Hutton JL, Pharaoh POD, Cooke RWI, Stevenson RC. Differential effects of pre-term birth and small gestational age on cognitive and motor development. *Arch Dis Child Fetal Neonatal Ed.* 1997; 76, F75–F81.
- Richards M, Hardy R, Kuh D, Wadsworth ME. Birth weight and cognitive function in the British 1946 birth cohort: longitudinal population based study. *Br Med J.* 2001; 322, 199–203.
- Orchinik LJ, Taylor HG, Espy KA, et al. Cognitive outcomes for extremely preterm/extremely low birth weight children in kindergarten. *J Int Neuropsychol Soc.* 2011; 17, 1067–1079.
- Richards M, Hardy R, Kuh D, Wadsworth ME. Birthweight, postnatal growth and cognitive function in a national UK birth cohort. *Int J Epidemiol.* 2002; 31, 342–348.
- Løhaugen GC, Gramstad A, Evensen KA, et al. Cognitive profile in young adults born preterm at very low birthweight. *Dev Med Child Neurol.* 2010; 52, 1133–1138.
- Strang-Karlsson S, Andersson S, Paile-Hyvärinen M, et al. Slower reaction times and impaired learning in young adults with birth weight <1500 g. *Pediatrics.* 2010; 125, e74–e82.
- Pyhälä R, Lahti J, Heinonen K, et al. Neurocognitive abilities in young adults with very low birth weight. *Neurology.* 2011; 77, 2052–2060.
- Shenkin SD, Starr JM, Deary IJ. Birth weight and cognitive ability in childhood: a systematic review. *Psychol Bull.* 2004; 130, 989–1030.
- Räikkönen K, Forsén T, Henriksson M, et al. Growth trajectories and intellectual abilities in young adulthood: the Helsinki Birth Cohort study. *Am J Epidemiol.* 2009; 170, 447–455.
- Broekman BF, Chan YH, Chong YS, et al. The influence of birth size on intelligence in healthy children. *Pediatrics.* 2009; 123, e1011–e1016.
- Eriksen W, Sundet JM, Tambs K. Birth weight standardized to gestational age and intelligence in young adulthood: a register-based birth cohort study of male siblings. *Am J Epidemiol.* 2010; 172, 530–536.
- Erickson K, Kritz-Silverstein D, Wingard DL, Barrett-Connor E. Birth weight and cognitive performance in older women: the Rancho Bernardo study. *Arch Womens Ment Health.* 2010; 13, 141–146.
- Shenkin SD, Starr JM, Pattie A, et al. Birth weight and cognitive function at age 11 years: The Scottish Mental Survey 1932. *Arch Dis Child.* 2001; 85, 189–197.
- Pearce MS, Deary IJ, Young AH, Parker L. Growth in early life and childhood IQ at age 11 years: the Newcastle Thousand Families Study. *Int J Epidemiol.* 2005; 34, 673–677.
- Costa AJ, Kale PL, Luiz RR, et al. Association between birthweight and cognitive function in middle age: the atherosclerosis risk in communities study. *Ann Epidemiol.* 2011; 21, 851–856.
- Tuvemo T, Jonsson B, Persson I. Intellectual and physical performance and morbidity in relation to height in a cohort of 18-year-old Swedish conscripts. *Horm Res.* 1999; 52, 186–191.
- Starr JM, Kilgour A, Pattie A, et al. Height and intelligence in the Lothian Birth Cohort 1921: a longitudinal study. *Age Ageing.* 2010; 39, 272–275.
- Laitala VS, Hjelmborg J, Koskenvuo M, et al. Shorter adult stature increases the impact of risk factors for cognitive impairment: a comparison of two Nordic twin cohorts. *Twin Res Hum Genet.* 2011; 14, 544–552.
- Daniels MC, Adair LS. Growth in young Filipino children predicts schooling trajectories through high school. *J Nutr.* 2004; 134, 1439–1446.
- Smith E, Hay P, Campbell L, Trollor JN. A review of the association between obesity and cognitive function across the lifespan: implications for novel approaches to prevention and treatment. *Obesity Rev.* 2011; 12, 740–755.
- Liu J, Raine A, Venables PH, Dalais C, Mednick SA. Malnutrition at age 3 years and lower cognitive ability at age 11 years: independence from psychosocial adversity. *Arch Pediatr Adolesc Med.* 2003; 157, 593–600.
- Australian Bureau of Statistics and Australian Institute of Health and Welfare. *The Health and Welfare of Australia's Aboriginal and Torres Strait Islander peoples. ABS cat. no. 4704.0, AIHW cat. no. IHW 11, 2003.* ABS: Canberra.
- Sayers SM, Mackerras D, Singh G, et al. An Australian aboriginal birth cohort: a unique resource for a life course study of an Indigenous population. A study protocol. *BMC Int Health Hum Rights.* 2003; 3, 1.
- Dubowitz LMS, Dubowitz V, Goldberg CA. *Clinical Manual: Gestational Age of the Newborn, 1977.* Addison-Wesley: London.
- Sayers SM, Powers JR. An evaluation of three methods used to assess gestational age of Aboriginal neonates. *J Paediatr Child Health.* 1992; 28, 312–317.
- Sayers S, Mackerras D, Halpin S, Singh G. Growth outcomes for Australian Aboriginal children aged 11 years who were born with intrauterine growth retardation at term gestation. *Paediatr Perinat Epidemiol.* 2007; 21, 411–417.
- Kuczmarski RJ, Ogden CL, Grummer-Strawn LM, et al. CDC growth charts: United States. *Advance Data.* 2000; 314, 1–27.
- Cairney S, Maruff P. Computerised tests of brain function for use with Indigenous people. In *Information Technology and Indigenous People* (eds. Dyson LE, Hendriks M, Grant S), 2007; pp. 257–259. Information Science Publishing: London.
- Dingwall KM, Lewis MS, Maruff P, Cairney S. Assessing cognition following petrol sniffing for Indigenous Australians. *Aust N Z J Psychiatry.* 2010; 44, 631–639.
- Keller H, Ayub BV, Saigal S, Bar-Or O. Neuromotor ability in 5- to 7-year-old children with very low or extremely low birthweight. *Dev Med Child Neurol.* 1998; 40, 661–666.

34. Somerfelt K, Andersson HW, Sonnander K, *et al.* Cognitive development of term small for gestational age children at five years of age. *Arch Dis Child.* 2000; 83, 25–30.
35. Jefferis BJ, Power C, Hertzman C. Birth weight, childhood socioeconomic environment, and cognitive development in the 1958 British birth cohort study. *Br Med J.* 2002; 325, 305–310.
36. Gorman BK. Birth weight and cognitive development in adolescence: causal relationship or social selection? *Soc Biol.* 2002; 49, 13–34.
37. Oddy WH, Kendall GE, Blair E, *et al.* Breast feeding and cognitive development in childhood: a prospective birth cohort study. *Paediatr Perinatal Epidemiol.* 2003; 17, 81–90.
38. National Health and Medical Research Council. *Nutrition in Aboriginal and Torres Strait Islander Peoples. An Information Paper,* 2000. Commonwealth of Australia: Canberra.
39. Canadian Paediatric Society. Growth assessment in Aboriginal children: is there need for change? *Paediatr Child Health.* 2004; 9, 477–479.